

Iorio's "high-precision measurement" of frame-dragging with the Mars Global Surveyor

Kris Krogh

Neuroscience Research Institute

University of California, Santa Barbara, CA 93106, USA

email: k_krogh@lifesci.ucsb.edu

November 10, 2007

Abstract

In two analyses of orbital data from the Mars Global Surveyor spacecraft, Iorio [1, 2] has claimed confirmation of the frame-dragging effect predicted by general relativity. Initially to an accuracy of 6%, and now 0.5%, exceeding the expected accuracy of NASA's Gravity Probe B. It is shown his results come from misinterpreting the MGS data and then altering a key time period.

1 Introduction

The Mars Global Surveyor (MGS) reached Mars in September 1997. While the main mission was imaging and mapping its surface, its magnetic and gravitational fields were also mapped. The latter began after aerobraking in the Mars atmosphere put the probe in an approximately circular polar orbit, at an altitude of 400 km, in February 1999. The gravitational field data were obtained by observing its orbital motion, via line-of-sight microwave Doppler and range measurements between the probe and Earth stations.

As first described by Lens and Thirring [3], general relativity predicts a rotating massive body would cause a rotational dragging of nearby space-time. In two recent papers, Iorio [1, 2] argues this subtle effect can be detected in the MGS orbit. Previous measurements have been reported by Ciufolini and coauthors [4, 5], using the Earth-orbiting LAGEOS and LAGEOS II satellites. In that case, the Lens-Thirring effect tends to be obscured by much larger ones, such as those of Earth and ocean tides, and the accuracy of those results has been questioned [6, 7].

A definitive finding is expected by year's end, with the announcement of results from NASA's Gravity Probe B. That experiment is an Earth-orbiting gyroscope, designed specifically to measure Lense-Thirring precession, to an accuracy of 1%. Iorio asserts he has already achieved a measurement, first to 6% and now 0.5% accuracy. This has been well publicized. (For example, see [8].) Here we will review the methods and calculations used to arrive at this extraordinary claim.

2 Analysis of the MGS Orbital Data

Iorio's Mars Global Surveyor papers [1, 2] are based on analysis of the MGS orbit by Konopliv *et al.* [9] Discussing their data, he writes in the first paper:

Here we are interested in particular in the out-of-plane portion of its orbit. The root-sum-square out-of-plane residuals over a five-year time interval spanning from 10 February 2000 to 14 January 2005 are shown in figure 1. They have been determined in a Mars-centered coordinate system. . . Due to improved modelling (orientation, gravity, angular momentum wheel de-saturations, and atmospheric drag), the average of such residuals amounts to 1.6 m.

This result can be well explained with the action of the gravitomagnetic field of Mars on the orbit of MGS.

The figure Iorio is referring to was reproduced from Konopliv *et al.* The original is shown as Figure 1 here. (Unlike Iorio's version, it shows six years of data.)

To compare the average residual error in the MGS orbit with the action of the gravitomagnetic field of Mars, Iorio starts with the gravitomagnetic Lens-Thirring precession of a satellite orbit due a rotating central body:

$$\dot{\Omega}_{LT} = \frac{2 G J}{c^2 a^3 (1 - e^2)^{3/2}} \quad (1)$$

(See Ciufolini and Wheeler [10] for a derivation.) $\dot{\Omega}_{\text{LT}}$ refers to the secular rate of change in the longitude of the orbit’s ascending node, G is the gravitational constant, J the rotating body’s angular momentum, c the speed of light, a and e are the orbit’s semi-major axis and eccentricity.

Referencing Christodoulidis *et al.* [11], he finds that a change in Ω_{LT} can be expressed as change in the position of the orbit’s ascending node N_{LT} as

$$\Delta N_{\text{LT}} = a \sqrt{1 + \frac{e^2}{2}} \sin i \Delta \Omega_{\text{LT}} \quad (2)$$

where i is the orbital plane’s inclination with respect to the central body’s equatorial plane. With ΔP representing a time interval, he combines these expressions to get

$$\langle \Delta N_{\text{LT}} \rangle = \frac{G J \Delta P \sin i \sqrt{1 + \frac{e^2}{2}}}{c^2 a^2 (1 - e^2)^{3/2}} \quad (3)$$

The factor of 2 in Eq. (1) is lost because $\langle \Delta N_{\text{LT}} \rangle$ represents the average change in position of the ascending node over this time interval.

The result from this equation is compared in both papers to the 1.6 meter out-of-plane residual orbit error from Konopliv *et al.* The parameter values he uses are the same in both cases, except for a J which increases slightly, and the values for ΔP , which change from 5 years to 5 years and two months. In his first paper, he gets a predicted 1.5 meter average Thirring precession of the ascending node, which he notes is within 6% of the value from Konopliv *et al.*

Instead of meters, the result of the second paper is presented as a ratio

$$\mu_{\text{meas}} \equiv \left. \frac{\langle \Delta N_{\text{res}} \rangle}{\langle \Delta N_{\text{LT}} \rangle} \right|_{\Delta P} = 0.9937 \pm 0.0053 \quad (4)$$

where the numerator refers to the 1.6 meter residual from Konopliv *et al.* The predicted Lens-Thirring precession, in the denominator, is 1.610 in this case. Of the uncertainty on the right, he attributes ± 0.0052 to uncertainty in the Mars angular momentum J , ± 0.0001 to uncertainty in the constant G , and *none* to the residual $\langle \Delta N_{\text{res}} \rangle$ from observations of the MGS trajectory.

What information is actually shown in Figure 1 from Konopliv *et al.*? The number of data points is related to a spacecraft trajectory model generated by the JPL Orbit Determination Program [12]. While data from the entire mission is processed to give a global best fit, the output is a sequence of separate trajectory segments, “data arcs,” 4 to 6 (Earth) days in length. Each plotted datum point corresponds to the intersection of two data arcs.

Ideally, the modeled segments would fit together as a smooth, continuous trajectory. Since the modeling cannot be perfect, there are mismatches. The orbital period of MGS is 2 hours, so a single data arc consists of many orbits. As Konopliv *et al.* [9] describe, to gauge and minimize mismatches between successive data arcs, each is extended to overlap the next by a full orbit. The RMS difference in the spacecraft positions is then computed over that orbit. (Iorio refers to these as root-sum-square residuals, although RMS indicates root-mean-square.)




Figure 1: RMS orbit overlap differences from Konopliv *et al* [9]. Reprinted with permission from Elsevier.

These position differences are taken in three directions: radial, along-track (the direction the spacecraft is moving), and normal to the orbit plane. Konopliv *et al.* plot these residual modeling errors in a three-part figure. Part (c), the “normal orbit difference,” is the basis of Iorio’s analysis, and is included above. The original point made was that, as modeling of the Mars gravity field and various perturbations have improved, the errors have declined. In this case, the normal average orbit error has decreased from a previous 3 to 1.6 meters.

This does *not* suggest a net precession of the orbit averaging 1.6 meters over the duration of the mission (ΔP). It refers to the average mismatch between successive data arcs. *Over the complete trajectory, the underlying directed errors might cancel, or accumulate to a large number of meters.* Even individually, the values do not specify orbital precessions.

The normal mismatch between overlapped orbits can be treated as having two parts: an angular tilt of the orbit plane around some axis through the Mars center of mass, plus a normal translation of the orbit plane. (The latter effect would not be expected in a rotationally symmetric system. For a planet with an irregular gravitational field, it may be significant.) Alone, a 1 meter normal translation of the orbit would give a 1 meter RMS difference, with no angular precession of the orbit.

Assuming the normal orbit difference is only due to Lense-Thirring precession, there would be no translation, and the corresponding orbital tilt would be exactly around the Mars rotation axis. In this case, the overlapped orbits would cross near the poles and separate at the Mars equatorial plane. If the ascending node positions differ there by 1 meter, the resulting RMS residual will be less, still not corresponding to what is assumed in Iorio’s calculation. Also, these numbers have no sign to indicate a direction of precession.

Moreover, there is no reason to expect the value 1.6 is unique. Lemoine *et al.* [13] and Yaun *et al.* [14] describe repeated decreases in the MGS orbit residuals as the accuracy of Mars gravity maps and perturbation modeling have improved. Nowhere in Konopliv *et al.* is it suggested that the present 1.6 m normal average orbit error could not be reduced by further refinement of the existing orbit model.

Another crucial element of Iorio’s analysis is his choice of the time period ΔP . According to him, the out-of-plane orbit error corresponds to a cumulative precession, increasing with time. The 1.6 m error from Konopliv *et al.* is based on 6 years of data, shown in Figure 1. Without explanation, Iorio’s first paper takes the same error to apply to a 5-year period. (His figure omits the data prior to 2000 to agree with that.)

In the second paper, he uses a ΔP equal to 5 years + 2 months, and obtains an order of magnitude better agreement between general relativity and experiment. This is possible because, while changing ΔP and with it the predicted error, he compares that to the same 1.6 m measured error once again. This time he notes

By using a time span ΔP of slightly more than 5 years (14 November 1999–14 January 2005), so to remove the first months of the mission more affected by orbital maneuvers and non-gravitational pertur-

bations, it is possible to obtain

$$\langle \Delta N_{\text{res}} \rangle = 1.6 \text{ m} \quad (5)$$

Let us, now, compare the result of [Eq. (5)] with the mean gravitomagnetic shift predicted by [Eq. (3)] over the same time span. . .

Two statistical tests are then used to show it is unlikely the close correspondence between his predicted value and that observed is due to chance. (This conclusion seems justified.) Stressing the objectivity of his findings, he also writes

Another important remark concerning our measurement is the following one: the interpretation of the MGS residuals presented in this paper and their production from the collected data were made by distinct and independent subjects in different times; moreover, the authors of [9] had not the Lense-Thirring effect in mind at all when they performed their analysis. Thus, the possibility of any sort of (un)conscious gimmicks, tricks or tweaking the data because the expected answer was known in advance [15] is a priori excluded in this test.

However, his results come from misinterpreting the MGS data, and from his own values of ΔP – twice altered in a way which gives closer agreement with general relativity.

3 Discussion

It would be surprising if an accurate test of Lens-Thirring precession (or any test of it) could be obtained from the Mars Global Surveyor trajectory. Unlike some spacecraft, it was not designed for measurement of its undisturbed inertial motion. That was possible, for example, with the Pioneer 10 and 11 space probes [16]. Those encountered no planetary atmospheres, and were spin-stabilized, such that frequent use of thrusters was not required.

In this case, AMD (angular momentum wheel desaturation) thrusters were fired 4 or 5 times per day until September 2001, when that was reduced to 1 or 2. Velocity uncertainties due to imprecision of the thrust are partially corrected from observations of the spacecraft motion, but that is limited by the one-dimensional nature of the Doppler velocity measurements.

From Konopliv *et al.*: “Normal orbit error is the greatest, and along-track is the least when the orbit plane is near edge-on as viewed from Earth (e.g., January 2001). For this geometry, the signature in the Doppler (which measures the velocity of the spacecraft in the Earth-Mars direction) is greatest for motion in the orbit plane, but is minimal for motion normal to the orbit plane.”

Referring to Figure 1, note the dips in the radial and along-track differences centered around January 2001, and the corresponding large hump in the normal orbit difference, where it increases by a factor of ten. This obviously contributes to the 1.6 m average normal orbit difference. We could agree with with Konopliv

et al. that this is due to *measurement* error. Certainly it cannot be attributed to a larger general relativistic precession occurring at that time.

Iorio's second paper states his value of ΔP was chosen "so to remove the first months of the mission more affected by orbital maneuvers and non-gravitational perturbations." However, there was no change in orbital maneuvers which would justify a time period beginning in November 1999. Also, Konopliv *et al.* point out that noise in the Doppler velocity signal was *least* during the first one and three quarters months, referred to as the "Gravity Calibration Orbit." This preceded deployment of the large high-gain antenna, which increased atmospheric drag.

At the relatively low altitude of MGS, the Martian atmosphere is a significant factor. From the spacecraft's asymmetric shape, drag due to its forward motion would be expected to cause out-of-plane forces. For a satellite in polar orbit, rotation of the upper atmosphere would also tend to produce a force causing an orbital precession. Solar heating of the spacecraft on emergence from the Mars shadow may be an additional factor.

None of the possible errors due to these or other orbital perturbations are estimated by Iorio, or included in his calculations. Following initial publication of this review on arXiv, another by Sindoni, Paris and Ialongo [17] has found the uncertainty of the orbit's precession exceeds that claimed in Iorio's second paper by at least a factor of 10000. Also see Felici [18].

4 Conclusions

Iorio makes the hypothesis that a certain measure of the error in the modeled Mars Global Surveyor orbit is due to the Lense-Thirring precession from general relativity. He assumes this measure is *precisely* 1.6 meters. This is shown to equal general relativity's prediction, within a factor of 0.9937, and with an uncertainty of ± 0.0053 . However, close agreement was obtained by calculating the predicted precession for a different time period than the actual period of observations.

Also, where he assumes 1.6 meters represents the average of a cumulative effect arising over 5 or 6 years, it actually characterizes an average RMS orbital mismatch arising over two "data arcs" – with a total span in the range of 8 to 12 days. The effects are generally not cumulative. Finally, this quantity is not a specific measure of movement of the orbit's ascending node, as assumed in Iorio's calculations.

Other approaches to measuring the Lens-Thirring effect are promising. In addition to the pending result from Gravity Probe B, a new experiment has been proposed by Turyshev *et al.* [19] This would observe gravitational bending of laser light by the Sun, using two small, relatively inexpensive satellites positioned on the far side of the Sun, and the International Space Station. Second-order bending would be measured for the first time, as well as that due to frame dragging caused by the Sun's rotation. The latter would be measured to 1% accuracy.

Acknowledgments

I am grateful to Stanley Robertson for first pointing out possible effects of the Mars atmosphere on the MGS orbit, and for helpful discussions. I thank Alexander Konopliv and E Myles Standish, for confirming my description of the MGS orbit error residuals, and for their support. Also the anonymous referees, for several useful suggestions.

References

- [1] Iorio L 2006 A note on the evidence of the gravitomagnetic field of Mars *Class. Quantum Grav.* **23** 5451 (*Preprint* gr-qc/0606092)
- [2] Iorio L 2007 High-precision measurement of frame-dragging with the Mars Global Surveyor spacecraft in the gravitational field of Mars *Preprint* gr-qc/0701042 versions v1-v5. (Note the unattributed changes in v6, following initial posting of this paper to arXiv: “root-sum-square” becomes “RMS,” residuals are now described as “overlap differences,” and the precise quantity 1.6 m becomes 1.613 m – giving even closer agreement with general relativity.)
- [3] Lense J and Thirring H 1918 Über den einfluss der eigenrotation der zentralkörper auf die bewegung der planeten und monde nach der Einsteinschen gravitationstheorie *Phys. Z.* **19** 156
- [4] Ciufolini I *et al.* 1998 Test of General Relativity and Measurement of the Lense-Thirring Effect with Two Earth Satellites *Science* **279** 2100
- [5] Ciufolini I and Pavlis E C 2004 A confirmation of the general relativistic prediction of the Lense-Thirring effect *Nature* **431** 958
- [6] Ries J C, Eanes R J and Tapley B D 2003 Lense-Thirring Precession Determination from Laser Ranging to Artificial Satellites *Nonlinear Gravitodynamics* ed R Ruffini and C Sigismondi (Singapore: World Scientific) pp 201-211
- [7] Iorio L 2005 On the reliability of the so far performed tests for measuring the Lense-Thirring effect with the LAGEOS satellites *New Astron.* **10** 603 (*Preprint* gr-qc/0411024)
- [8] Clark S 2007 Loner stakes claim to gravity prize *New Scientist* **2587** 14
- [9] Konopliv A S, Yoder C F, Standish E M, Yuan D N, Sjogren W L 2006 A global solution for the Mars static and seasonal gravity, Mars orientation, Phobos and Deimos masses, and Mars ephemeris *Icarus* **182** 23
- [10] Ciufolini I and Wheeler J A 1995 *Gravitation and Inertia* (Princeton: Princeton University Press)

- [11] Christodoulidis D C *et al.* 1988 Observed tidal braking in the Earth/Moon/Sun system *J. Geophys. Res.* **93** 6216
- [12] Moyer T D 2003 *Formulation for Observed and Computed Values of Deep Space Network Data Types for Navigation* (Hoboken: Wiley-Interscience)
- [13] Lemoine F G *et al.* 2001 An improved solution of the gravity field of Mars (GMM-2B) from Mars Global Surveyor *J. Geophys. Res.* **106** 23359
- [14] Yuan D N *et al.* 2001 Gravity field of Mars: a 75th Degree and Order Model *J. Geophys. Res.* **106** 23377
- [15] Weinberg S 1993 *Dreams of a final theory* (New York: Vintage Books)
- [16] Anderson J D *et al.* 2002 Study of the anomalous acceleration of Pioneer 10 and 11 *Phys. Rev. D* **65** 082004 (*Preprint* gr-qc/0104064)
- [17] Sindoni G, Paris C and Ialongo P 2007 On the Systematic Errors in the Detection of the Lense-Thirring Effect with a Mars Orbiter *Preprint* gr-qc/0701141
- [18] Felici G 2007 The meaning of systematic errors, a comment to “Reply to On the Systematic Errors in the Detection of the Lense-Thirring Effect with a Mars Orbiter,” by Lorenzo Iorio *Preprint* gr-qc/0703020
- [19] Turyshev S G *et al.* 2005 Fundamental physics with the laser astrometric test of general relativity *ESA Spec. Publ.* **588** 11 (*Preprint* gr-qc/0506104)